



CONTENTS *continued*

	<i>Page</i>
Typhoid fever epidemic following a wedding reception . . . <i>Charles T. Caraway and John M. Bruce</i>	427
Public health services at the 1960 Winter Olympics <i>Richard F. White, Malcolm H. Merrill, and William W. Stiles</i>	431
Measuring social restoration performance of public psychi- atric hospitals <i>Charles E. Rice, David G. Berger, Lee G. Sewall, and Paul V. Lemkau</i>	437
The USSR test as a screening test in a public health labora- tory <i>Daniel Widelock, Joseph Portnoy, John Truelove, Anna D. Reynolds, and Jules Vandow</i>	447
Detecting viruses on nonporous surfaces by use of cotton swab technique <i>Kenneth Keller, Herschul Braspeninckx, and Morton Kasdan</i>	453
Short reports and announcements:	
Municipal water pollution control needs	377
Housing standards. Legal note	378
Zoonoses research center	384
Publication announcements	419
Guidelines for mental health service facilities	424
Child health center to be established	426
International mail pouch	430
Psychiatric case register	446
Accidental drownings at home. Epidemiological note	452
Wider use of research findings urged	458
Federal publications	459



MANAGING DIRECTOR

WILLIAM H. STEWART, M.D.
*Chief, Division of Public Health
Methods*

BOARD OF EDITORS

ERNEST L. STEBBINS, M.D., M.P.H.
Chairman

FRANCIS A. ARNOLD, JR., D.D.S.
A. L. CHAPMAN, M.D.

W. PALMER DEARING, M.D.

HERBERT R. DOMKE, M.D., DR.P.H.

ROBERT DYAR, M.D., DR.P.H.

WESLEY E. GILBERTSON, M.S.P.H.

ROGER W. HOWELL, M.D.

CHARLES V. KIDD, PH. D.

KARL M. MASON, B.S.S.E., M.P.H.

JAMES R. SHAW, M.D.

RUTH SLEEPER, R.N., M.A.

HELEN M. WALLACE, M.D.

STAFF

Marcus Rosenblum *Executive Editor*

Winona Carson *Managing Editor*

Martha Seaman *Asst. Managing Editor*

Eugene Fite *Art Editor*

Address correspondence to Executive Editor

Opinions expressed are the authors' and do not necessarily reflect the views of *Public Health Reports* or the Public Health Service. Trade names are used for identification only and do not represent an endorsement by the Public Health Service.

U.S. DEPARTMENT OF HEALTH, EDUCATION, AND WELFARE

ABRAHAM RIBICOFF, *Secretary*

PUBLIC HEALTH SERVICE

LUTHER L. TERRY, *Surgeon General*

PHR

S
Y
N
O
P
S
E
S

HENNESSY, ALBERT V. (University of Michigan), and **DAVENPORT, FRED M.:** *Relative merits of aqueous and adjuvant influenza vaccines when used in a two-dose schedule. Public Health Reports, Vol. 76, May 1961, pp. 411-419.*

Broad and high levels of antibody against all known families of influenza A virus may be induced in both children and adults either by two doses of adjuvant polyvalent vaccine or by aqueous polyvalent vaccine followed by adjuvant polyvalent vaccine. Antibodies so produced remain at high levels for a long period of time while antibodies produced

by two doses of aqueous vaccine fall to low levels in 1 year.

In children, two very small doses of virus in adjuvant vaccine gave high levels of antibody even when the interval between inoculations was only 8 weeks.

The findings reported indicate that high, broad, and persistent antibody levels can be achieved in man with a minimum of inoculations and materials.

RICE, CHARLES E. (Veterans Administration Hospital, Perry Point, Md.), **BERGER, DAVID G., SEWALL, LEE G., and LEMKAU, PAUL V.:** *Measuring social restoration performance of public psychiatric hospitals. Public Health Reports, Vol. 76, May 1961, pp. 437-446.*

Interest in the problem of how to evaluate the performance of public psychiatric hospitals is growing. The Medical Audit Plan for Psychiatric Hospitals is a research program organized to develop methods for appraising mental hospital effectiveness with respect to the achievement of certain hospital goals or objectives.

This paper describes a method for measuring psychiatric hospital performance in the area of one such goal, social restoration. A hospital's performance in socially restoring its patients may be evaluated at each of three "levels": the extent to which the hospital authorizes

the release of patients to the community; the extent to which such patients remain in the community following their release; and the extent to which released patients adjust to living in the community.

The measurement of these levels, based on data obtained by studying cohorts of psychiatric patients, is described in detail. Data gathering methods are described briefly. These methods, should they prove workable, will provide a basis for evaluation of the effectiveness of public psychiatric hospitals and a guide for administrative policy and program decisions.

PHR

S
Y
N
O
P
S
E
S

WIDELOCK, DANIEL (New York City Department of Health); **PORTNOY, JOSEPH**; **TRUELOVE, JOHN**; **REYNOLDS, ANNA D.**; and **VANDOW, JULES**: *The USR test as a screening test in a public health laboratory. Public Health Reports, Vol. 76, May 1961, pp. 447-452.*

At the New York City Department of Health a serologic study was undertaken in an attempt to cut the mounting costs of syphilis serology and to solve the problem of the biologic false positive reaction.

During the study 100,000 blood specimens were tested simultaneously by the unheated serum reagin (USR) and VDRL slide tests. All serums reactive with the USR or VDRL tests were further tested by the Kolmer complement fixation test. The findings indicate that the USR test may be employed as a screening procedure in a public health laboratory with a saving in space, personnel, and time.

Treponemal tests, the Reiter protein complement fixation (RPCF) test and the *Treponema pallidum* complement fixation 50 (tpcf-50) test were carried out on

1,885 serums positive by the USR or VDRL tests, or by both, for which a clinical diagnosis was available. Results indicated that the RPCF test may be substituted for the more expensive tpcf-50 test.

A public health laboratory procedure may consist of the following:

1. Use of the USR test as a screening procedure, with all negative results being reported immediately and all USR reactors being tested further.

2. Testing of positive reactors by VDRL titrations, Kolmer complement fixation test, and RPCF test.

3. If additional testing is required, employment of the Nichols strain of *T. pallidum* in an approved test, the fluorescent treponemal antibody (FTA) test, for example.

KELLER, KENNETH; **BRASPENNINCKX, HERSCHUL**; and **KASDAN, MORTON** (University of Louisville School of Medicine): *Detecting viruses on nonporous surfaces by use of the cotton swab technique. Public Health Reports, Vol. 76, May 1961, pp. 453-458.*

A preliminary study was performed to determine whether the standard cotton swab-rinse technique for detecting bacterial contaminants could also be used for the recovery of viruses from nonporous surfaces. The results indicated that, under the experimental conditions described, the percentage of recovery of bacteriophage T1 virus approximates the precision obtained in detecting bacterial contaminants.

The data show that the cotton swab-rinse method is sensitive enough to detect virus when as few as 16 particles are initially applied to a 4-square-inch test surface area and air dried for 1 hour.

As few as 12 TCD₅₀, the virus dose which gives rise to cytopathogenic changes in 50 percent of the tissue culture tubes, of poliovirus 1 per 4 square inches were detected, using five HeLa cell culture tubes per sample, when sampling was done 5 minutes after the test surfaces were contaminated.

When the test surfaces were contaminated and then allowed to air-dry for 1 hour, the end point of poliovirus detection occurred between contaminating inoculum titers of between 6,310 TCD₅₀ and 631 TCD₅₀ per 4 square inches.

Information for Contributors

PUBLIC HEALTH REPORTS welcomes from any source all contributions of value to public health.

Most of the readers of *Public Health Reports* are practicing public health officials. About 10 percent of the monthly circulation of *Public Health Reports* goes overseas. About half of the domestic circulation goes to Federal, State, and local government agencies concerned with health and related health interests. A quarter goes to institutions accredited for teaching in health and related fields, to teachers, and to libraries. The journal also reaches research institutions, hospitals, and professional and voluntary public health organizations.

Tearsheets. In lieu of reprints, senior authors are provided with 50 to 100 sets of tearsheets after publication. Associate authors receive a smaller number.

Manuscript review. Manuscripts submitted for publication are reviewed by technical experts, and authors are given the benefit of their comments before type is set. Authors also receive edited typescripts for approval and are given the opportunity to correct galley proofs. Authors are responsible for the accuracy and validity of all material, including tables, charts, and references. Special editorial assistance in preparing or revising manuscripts is available on request, to the limit of staff resources.

Manuscripts are reviewed with the understanding that they have not been committed for publication elsewhere. Appropriate information should be provided if a paper has been given or is prepared for presentation at a meeting.

Manuscript form. Authors will facilitate review and publication if they submit an original and three carbon copies of their manuscripts. All copy should be typed double spaced, and each page should end with a completed paragraph. Of course, several paragraphs may appear on a typed page.

References should be given in the style used by *Public Health Reports*.

Footnotes should be worked into the text or offered as supplemental items.

Authors are expected to recognize scientific contributions by those who have assisted in their papers only if such contributions warrant mention in the text or in the paragraph identifying the authors. It is not the policy of *Public Health Reports* to publish "acknowledgments."

Synopses. Authors are requested to provide a 200-word synopsis of appropriate papers. The staff will supply on request information offering guidance on the preparation of synopses.

Index listings. *Public Health Reports* is listed in the annual *Cumulated Index Medicus* (American Medical Association), in the monthly *Index Medicus* (National Library of Medicine), in the *Engineering Index*, and in the *Hospital Literature Index*.

Bound copies. Librarians and others should preserve their copies for binding, as the Public Health Service does not supply bound copies. Indexes are published each year in the December issue.

PUBLIC HEALTH MONOGRAPHS, edited and issued by *Public Health Reports*, must be submitted through constituent agencies of the Department of Health, Education, and Welfare.

Most Public Health Monographs are placed on sale by the Superintendent of Documents; series subscriptions are not available. Monographs are not included in subscriptions to *Public Health Reports*.

Address correspondence on editorial matters to: Executive Editor, Public Health Reports, Public Health Service, U.S. Department of Health, Education, and Welfare, Washington 25, D.C.